arXiv:1101.3614v1 [math.CO] 19 Jan 2011

Direct bijective computation of the generating series for 2 and 3-connection coefficients of the symmetric group

Alejandro H. Morales and Ekaterina A. Vassilieva

January 20, 2011

Abstract

We evaluate combinatorially certain connection coefficients of the symmetric group that count the number of factorizations of a long cycle as a product of three permutations. Such factorizations admit an important topological interpretation in terms of unicellular constellations on orientable surfaces. Bijective computations of these coefficients are so far limited to very special cases. Thanks to a new bijection that refines the work of Schaeffer and Vassilieva in [16] and [17], we give an explicit closed form evaluation of the generating series for these coefficients. The main ingredient in the bijection is a modified oriented tricolored tree tractable to enumerate. Finally, reducing this bijection to factorizations of a long cycle into two permutations, we get the analogue formula for the corresponding generating series.

Contents

1 Introduction **2**
1.1 Generating series for connection coefficients . . . 2
1.2 Background . . . 2
1.3 Outline of paper . . . 3

2 Combinatorial reformulation **3**
2.1 Cacti and partitioned cacti . . . 3
2.2 Thorn cactus trees . . . 6
2.3 Reformulation of the main theorem . . . 7

3 Description of the bijection **8**
3.1 Additional definitions . . . 8
3.2 Bijective mapping Θ for 3-cacti that preserves type . . . 9
3.2.1 The ordered thorn cactus tree $\tilde{\tau}$: . . . 9
3.2.2 The ordered set χ and the permutations σ_1 and σ_2 . . . 15
3.3 Showing the map Θ is a bijection . . . 15

4 Proof of Proposition 2: computation of the number of thorn cactus trees **16**

5 Proof of Corollary 1 and restriction of bijection Θ when $\nu = [1^n]$ **20**

1 Introduction

1.1 Generating series for connection coefficients

In what follows, we denote by $\lambda \vdash n$ an integer partition of n and $\ell(\lambda) = k$ the length or number of parts of λ.

Let S_n be the symmetric group on n elements, and $\mathcal{C}_\lambda$ be the conjugacy class in S_n of permutations with cycle type λ, where $\lambda \vdash n$. Given $\lambda^{(1)}, \lambda^{(2)}, \ldots, \lambda^{(r)}, \mu \vdash n$, let $k^{\mu}_{\lambda^{(1)},\ldots,\lambda^{(r)}}$ be the number of ordered factorizations in S_n of a fixed permutation $\gamma \in \mathcal{C}_\nu$ as a product $\alpha_1 \cdots \alpha_r$ of r permutations $\alpha_i \in \mathcal{C}_{\lambda^{(i)}}$. These numbers are called *connection coefficients* of the symmetric group. The problem of computing these coefficients has received significant attention and its history and references can be found in [9]. We focus on the cases $k^n_{\lambda,\mu}$ and $k^n_{\lambda,\mu,\nu}$: *i.e.* when $r = 2$ and 3, $\nu = [n]$ and γ is the long cycle $\gamma_n = (1, 2, \ldots, n)$.

In addition, for $\lambda \vdash n$, we use the monomial symmetric function $m_\lambda(\mathbf{x})$ which is the sum of all different monomials obtained by permuting the variables of $x_1^{\lambda_1} x_2^{\lambda_2} \cdots$, and the power symmetric function $p_\lambda(\mathbf{x})$, defined multiplicatively as $p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots$ where $p_n(\mathbf{x}) = m_n(\mathbf{x}) = \sum_i x_i^n$. Also, if $n_i(\lambda)$ is the number of parts of λ that are equal to i so that $\lambda = [1^{n_1(\lambda)}, 2^{n_2(\lambda)}, \ldots]$, let $Aut(\lambda) = \prod_i n_i(\lambda)!$.

Our combinatorial results can be stated as follows:

Theorem 1. *The numbers $k^n_{\lambda,\mu,\nu}$ of factorizations of the long cycle γ_n into an ordered product of three permutations of types $\lambda, \mu,$ and ν respectively satisfy:*

$$\sum_{\lambda,\mu,\nu \vdash n} k^n_{\lambda,\mu,\nu} p_\lambda(\mathbf{x}) p_\mu(\mathbf{y}) p_\nu(\mathbf{z}) = \sum_{\lambda,\mu,\nu \vdash n} \frac{n!^2 M^{(n-1)}_{\ell(\lambda),\ell(\mu),\ell(\nu)}}{\binom{n-1}{\ell(\lambda)-1}\binom{n-1}{\ell(\mu)-1}\binom{n-1}{\ell(\nu)-1}} m_\lambda(\mathbf{x}) m_\mu(\mathbf{y}) m_\nu(\mathbf{z}), \tag{1}$$

where

$$M^{(n-1)}_{\ell(\lambda),\ell(\mu),\ell(\nu)} = \binom{n-1}{\ell(\nu)-1} \sum_{g \geq 0} \binom{n-\ell(\mu)}{\ell(\lambda)-1-g} \binom{n-\ell(\nu)}{g} \binom{n-1-g}{n-\ell(\mu)}.$$

Corollary 1. *[14]*

$$\sum_{\lambda,\mu \vdash n} k^n_{\lambda,\mu} p_\lambda(\mathbf{x}) p_\mu(\mathbf{y}) = \sum_{\lambda,\mu \vdash n} \frac{n(n-\ell(\lambda))!(n-\ell(\mu))!}{(n+1-\ell(\lambda)-\ell(\mu))!} m_\lambda(\mathbf{x}) m_\mu(\mathbf{y}) \tag{2}$$

We will see in Section 2 that the coefficients on the right hand sides of (1) and (2) are non-negative integers.

Remark 1. *Equations* (1) *and* (2) *can be obtained algebraically using the irreducible characters of the symmetric group, the Murnaghan-Nakayama rule, and symmetric function identities (see [11]). Here, we derive these equations through bijections.*

1.2 Background

In the setting of the connection coefficients $k^n_{\lambda^{(1)},\cdots,\lambda^{(r)}}$, we define the *genus* $\mathsf{g}(\lambda^{(1)}, \ldots, \lambda^{(r)})$ of the partitions $\lambda^{(i)}$ by the equation

$$\ell(\lambda^{(1)}) + \cdots + \ell(\lambda^{(r)}) = (r-1)n + 1 - 2\mathsf{g}(\lambda^{(1)}, \cdots, \lambda^{(r)}).$$

We can take g to be a non-negative integer, since otherwise it is easy to show that $k^n_{\lambda^{(1)}\cdots\lambda^{(r)}} = 0$.

Regarding the evaluation of $k^n_{\lambda,\mu}$, using an inductive argument Bédard and Goupil [1] first found a formula for the case $\mathsf{g}(\lambda,\mu)=0$. This was extended by Goulden and Jackson [6] to evaluate $k^n_{\lambda^{(1)},\ldots,\lambda^{(r)}}$ in the case $\mathsf{g}(\lambda^{(1)},\ldots,\lambda^{(r)})=0$ via a bijection with a set of ordered rooted r-cacti on n r-gons. Later, using characters of the symmetric group and a combinatorial development, Goupil and Schaeffer [9] derived an expression for connection coefficients of arbitrary genus as a sum of positive terms (see Biane [3] for a succinct algebraic derivation; Poulalhon and Schaeffer [15] and Irving [10] for further generalizations). Recently, Chapuy [4] and [5] found a remarkable combinatorial identity for $\sum_{\lambda,\mu,\ell(\lambda)=a,\ell(\mu)=b} k^n_{\lambda,\mu}$ by identifying $2\mathsf{g}$ special elements in the cycles, called *trisections*, and slicing the cycles at these to obtain lower genus factorizations of the long cycle.

Interestingly, if the permutations are associated to set partitions that refine the partitions induced by the disjoint cycle decomposition (*i.e.* we color the cycles allowing repeated colors), then explicit direct bijections and closed expressions are attained (see Lass [13], Goulden and Nica [8], and Bernardi [2]). In this paper we follow this approach and introduce the notion of *partitioned tricolored (bicolored)* 3*-cacti (maps)* of given type, refining the work of Schaeffer and Vassilieva in [16] and [17], and use a purely combinatorial argument to derive the explicit generating series for $k^n_{\lambda,\mu,\nu}$ and $k^n_{\lambda,\mu}$ in Equations (1) and (2) respectively.

1.3 Outline of paper

The paper is organized as follows: in Section 2 we introduce the *partitioned* 3*-cacti* and the *thorn cactus trees* (the enumeration of the latter is postponed to Section 4) and relate them via a bijection Θ described in Section 3. Finally, in Section 5 we prove Corollary 1.

2 Combinatorial reformulation

2.1 Cacti and partitioned cacti

Factorizations in the symmetric group counted by $k^n_{\lambda,\mu,\nu}$ admit a direct interpretation in terms of 3*-cacti* (*i.e. unicellular* 3*-constellations*) with white, black, and grey vertices of respective degree distribution λ, μ, and ν. Within a topological point of view, 3-cacti are specific maps (graphs embedded in oriented surfaces defining a 2-cell decomposition of this oriented surface into a finite number of vertices, edges, and faces homeomorphic to open discs; see [12] for more details about maps and their applications) with *one* white face and n black faces, such that all the black faces are triangles that do not share edges. Each triangle is composed of exactly a white, a black, and a grey vertex following each other in clockwise order. Each white, black, or grey vertex corresponds to a cycle of α_1, α_2, or α_3 respectively, and the cycle is encoded by the local *counter-clockwise* order of the triangles the vertex belongs to.

The degree of a vertex in a cacti is the number of such triangles, and the degree distribution of the vertices of a given color is the partition of n formed by the degrees of all the vertices of this color. The fact that $\alpha_1\alpha_2\alpha_2=\gamma_n$ corresponds to saying that traversing the map starting on the white vertex of the triangle labelled 1 and keeping the unbounded face on the *right* we visit, in order, the white-black edges belonging to triangles $1,2,\ldots,n$. Figure 1 shows a 3-cactus embedded on the sphere (genus 0) and a 3-cactus embedded on a torus (genus 1).



Figure 1: Two examples of cacti embedded in a surface of genus 0 (left) and genus 1 (right)

Example 1. *The cactus on the left hand side in Figure 1 can be associated to the three permutations:*

$$\begin{aligned} \alpha_1 &= (1)(24)(3)(5) \quad (3) \\ \alpha_2 &= (1)(23)(45) \quad (4) \\ \alpha_3 &= (15)(2)(3)(4) \quad (5) \end{aligned}$$

The type of this cactus is $\lambda = [1^3, 2^1]$, $\mu = [1^1, 2^2]$, $\nu = [1^3, 2^1]$.
The cactus on the right hand side corresponds to the three permutations:

$$\begin{aligned} \alpha_1 &= (1236)(4)(5) \quad (6) \\ \alpha_2 &= (153)(2)(4)(6) \quad (7) \\ \alpha_3 &= (134)(2)(5)(6) \quad (8) \end{aligned}$$

The type of this cactus is $\lambda = [1^2, 4^1]$, $\mu = [1^3, 3^1]$, $\nu = [1^3, 3^1]$.

Partitioned 3-cacti

Cacti with a given vertex degree distribution are in general non-planar and non-recursive objects, and no direct bijection is known to compute their cardinality. However, we provide an interpretation of the formal power series $\sum_{\lambda,\mu,\nu\vdash n} k^n_{\lambda,\mu,\nu} p_\lambda(\mathbf{x})p_\mu(\mathbf{y})p_\nu(\mathbf{z})$ as a generating function for *partitioned cacti*. Introducing a new bijective mapping for partitioned cacti, we are able to give an explicit formula for this generating function.

To define partitioned cacti we use π to denote a set partition of $[n] = \{1, 2, \dots, n\}$ with blocks $\{\pi^1, \dots, \pi^p\}$. The type of a set partition, *type*$(\pi) \vdash n$, is the integer partition of n obtained by considering the cardinalities of the blocks of π. We are now ready for the definition:

Definition 1. *(Partitioned 3-cacti) For* λ, μ, $\nu \vdash n$, *we let* $\mathcal{C}(\lambda, \mu, \nu)$ *be the set of partitioned 3-cacti* $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2)$ *composed of set partitions* π_1,π_2, *and* π_3 *with respective type* λ, μ, *and* ν; *and permutations* $\alpha_1, \alpha_2 \in S_n$ *such that:*

- *each block of* π_1 *is the union of cycles of* α_1,
- *each block of* π_2 *is the union of cycles of* α_2,
- *each block of* π_3 *is the union of cycles of* $\alpha_3 = \alpha_2^{-1} \circ \alpha_1^{-1} \circ \gamma_n$.

The blocks of π_1, π_2, *and* π_3 *are denoted* $\pi_1^{(i)}, \pi_2^{(j)}, \pi_3^{(k)}$ *with the only restriction that* $1 \in \pi_1^{(\ell(\lambda))}$. *Let* $C(\lambda, \mu, \nu) = |\mathcal{C}(\lambda, \mu, \nu)|$.

Note that in terms of factorizations, partitioned cacti can be viewed as ordered factorizations of γ_n where we label the cycles of permutations, using distinct sets of labels for each permutation, allowing repeated labels within the permutation. Then the parts of λ are the number of elements in $[n]$ belonging to cycles of α_1 with the same label. An equivalent statement holds for μ and ν. From a topological point of view, partitioned cacti can be seen as 3-cacti encoded by the permutations α_1, α_2, and α_3 whose vertices of each color are partitioned. These partitions of the vertices are encoded by the set partitions of $[n]$ π_1, π_2, and π_3 respectively.

Example 2. *Take the cactus on the right hand side of Figure 1and add partitions* $\pi_1 = \{\pi_1^{(1)}, \pi_1^{(2)}\}, \pi_2 = \{\pi_2^{(1)}, \pi_2^{(2)}\}$, $\pi_3 = \{\pi_3^{(1)}, \pi_3^{(2)}, \pi_3^{(3)}\}$, *where* $\pi_1^{(1)} = \{4,5\}$, $\pi_1^{(2)} = \{1,2,3,6\}$; $\pi_2^{(1)} = \{1,3,4,5\}$, $\pi_2^{(2)} = \{2,6\}$; $\pi_3^{(1)} = \{1,3,4,6\}$, $\pi_3^{(2)} = \{2\}$, $\pi_3^{(3)} = \{5\}$. *This gives a partitioned cactus* $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) \in \mathcal{C}([2^1, 4^1], [2^1, 4^1], [1^2, 4^1])$ *depicted in Figure 2. Similarly to [16], we associate a particular shape to each of the blocks of the partitions.*



Figure 2: Example of a Partitioned 3-Cactus

Link between cacti and partitioned cacti

Consider the partial order on integer partitions given by refinement. That is $\lambda \preceq \mu$ if and only if the parts of μ are unions of parts of λ, and we say that μ is *coarser* than λ. If $\lambda \preceq \mu$ let $\overline{R}_{\lambda,\mu}$ be the number of ways to *coarse* λ to obtain μ. For example, if $\lambda = [1^2, 2^2]$ and $\mu = [1, 2, 3]$ then $\overline{R}_{\lambda,\mu} = 4$.

Remark 2. *If* $m = \ell(\lambda)$ *and* $p = \ell(\mu)$ *then* $\overline{R}_{\lambda,\mu}$ *is equal to the number of unordered partitions* $\pi = \{\pi^1, \dots, \pi^p\}$ *of the set* $[m]$ *such that* $\mu_j = \sum_{i \in \pi^j} \lambda_i$ *for* $1 \le j \le p$. *Therefore, for the monomial and power symmetric functions,* m_λ *and* p_λ, *we have:* $p_\lambda = \sum_{\mu \succeq \lambda} Aut(\mu) \overline{R}_{\lambda,\mu} m_\mu$ *[18, Prop.7.7.1].*

We use this partial order on integer partitions to obtain an immediate relation between $C(\lambda, \mu, \nu)$ and $k^n_{\lambda,\mu,\nu}$.

Proposition 1. *For partitions* $\rho, \delta, \epsilon \vdash n$ *we have :*

$$C(\rho, \delta, \epsilon) = \sum_{\lambda \preceq \rho, \mu \preceq \delta, \nu \preceq \epsilon} \overline{R}_{\lambda\rho} \overline{R}_{\mu\delta} \overline{R}_{\nu\epsilon} k^n_{\lambda,\mu,\nu}. \tag{9}$$

Proof. Let $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) \in \mathcal{C}(\rho, \delta, \epsilon)$. If $\alpha_1 \in \mathcal{C}_\lambda$, $\alpha_2 \in \mathcal{C}_\mu$, and $\alpha_3 = \alpha_2^{-1}\alpha_1^{-1}\gamma_n \in \mathcal{C}_\nu$ then by the definition of the partitioned cacti, we have that *type*$(\pi_1) = \rho \succeq \lambda$, *type*$(\pi_2) = \delta \succeq \mu$, and *type*$(\pi_3) = \epsilon \succeq \nu$. Thus, if we further refine $C(\rho, \delta, \epsilon)$ by the cycle types of the permutations, *i.e.* if

$$\mathcal{C}_{\lambda,\mu,\nu}(\rho, \delta, \epsilon) = \{(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) \in \mathcal{C}(\rho, \delta, \epsilon) \mid (\alpha_1, \alpha_2, \alpha_2^{-1}\alpha_1^{-1}\gamma_n) \in \mathcal{C}_\lambda \times \mathcal{C}_\mu \times \mathcal{C}_\nu\},$$

then $\mathcal{C}(\rho,\delta,\epsilon) = \bigcup_{\lambda\preceq\rho,\mu\preceq\delta,\nu\preceq\epsilon} \mathcal{C}_{\lambda,\mu,\nu}(\rho,\delta,\epsilon)$ where the union is disjoint. Finally, if $C_{\lambda,\mu,\nu}(\rho,\delta,\epsilon) = |\mathcal{C}_{\lambda,\mu,\nu}(\rho,\delta,\epsilon)|$ then it is easy to see that $C_{\lambda,\mu,\nu}(\rho,\delta,\epsilon) = \overline{R}_{\lambda\rho}\overline{R}_{\mu\delta}\overline{R}_{\nu\epsilon}k^n_{\lambda,\mu,\nu}$. □

Using $p_\lambda = \sum_{\rho\succeq\lambda} R_{\lambda\rho}m_\rho$ Proposition 1 is equivalent to:

$$\sum_{\lambda,\mu,\nu\vdash n} k^n_{\lambda,\mu,\nu} p_\lambda(\mathbf{x})p_\mu(\mathbf{y})p_\nu(\mathbf{z}) = \sum_{\lambda,\mu,\nu\vdash n} Aut(\lambda)Aut(\mu)Aut(\nu)\, C(\lambda,\mu,\nu)\, m_\lambda(\mathbf{x})m_\mu(\mathbf{y})m_\nu(\mathbf{z}). \tag{10}$$

Remark 3. *Let* $(\pi_1,\pi_2,\pi_3,\alpha_1,\alpha_2) \in \mathcal{C}(\rho,\delta,\epsilon)$ *with* $\ell(\rho)+\ell(\delta)+\ell(\epsilon) = 2n+1$*, and* $\alpha_3 = \alpha_2^{-1}\alpha_1^{-1}\gamma$*. If* $\alpha_1 \in \mathcal{C}_\lambda$*,* $\alpha_2 \in \mathcal{C}_\mu$*,* $\alpha_3 \in \mathcal{C}_\nu$*, then* $\ell(\lambda)+\ell(\mu)+\ell(\nu) = 2n+1-2\mathsf{g}(\lambda,\mu,\nu) \leq 2n+1$*. But* $\ell(\lambda) \geq \ell(\rho)$*,* $\ell(\mu) \geq \ell(\delta)$*, and* $\ell(\nu) \geq \ell(\epsilon)$ *therefore* $\lambda = \rho$*,* $\mu = \delta$*, and* $\nu = \epsilon$*; and* π_1*,* π_2*, and* π_3 *are the underlying set partitions in the cycle decompositions of* α_1, α_2*, and* α_3 *respectively. In this case,* $C(\lambda,\mu,\nu) = k^n_{\lambda,\mu,\nu}$ *(*$\mathsf{g}(\lambda,\mu,\nu) = 0$*) and this genus zero coefficient is* $\frac{n^2(\ell(\lambda)-1)!(\ell(\mu)-1)!(\ell(\nu)-1)!}{Aut(\lambda)Aut(\mu)Aut(\nu)}$ *as shown in [6, Thm. 2.2].*

As mentioned before, explicit computation of the right hand side of equation (10) is made possible thanks to a new bijective description of partitioned cacti of given type. Partitioned cacti are indeed in one-to-one correspondence with particular sets of thorn cactus trees (and three additional simple combinatorial objects), which are recursive planar objects one can compute with classical methods. Next we gradually define such trees.

2.2 Thorn cactus trees

Let $\mathcal{T}(p_1,p_2,p_3)$ be the set of rooted tricolored trees t with p_1 white vertices, p_2 black vertices, and p_3 grey vertices such that: (i) the root of t is a white vertex, (ii) a white vertex has black descendants, (iii) a black vertex has grey descendants, and (iv) a grey vertex has white descendants.

Given a tree $\mathsf{t} \in \mathcal{T}(p_1,p_2,p_3)$, take a vertex u and one of its descendants v. We allow adding an edge between u and *the rightmost* descendant of v, say vertex w. Whenever this happens, we say that there is a *triangle* rooted at u, and the additional edge $\{\mathsf{u},\mathsf{w}\}$ is called a *non-tree edge.* The vertex v, opposite to a non-tree edge, is called a *middle vertex.* The tree t with these triangles is called a *cactus tree.* More specifically:

Definition 2 (Cactus Tree)**.** *Let* $\mathcal{CT}(p_1,p_2,p_3,g,w,b)$ *be the set of rooted cactus trees* τ *consisting of an underlying tree* $\mathsf{t} \in \mathcal{T}(p_1,p_2,p_3)$ *and in addition* t *has:* g *triangles rooted in a grey vertex,* w *triangles rooted in a white vertex,* b *triangles rooted in a black vertex such that:*
(v) triangles are composed of a white, a black, and a grey vertex. The vertices are ordered clockwise cyclically as white-black-grey-white.

Lastly, we consider a related set of cactus trees with *thorns* or half edges on some of its vertices:

Definition 3 (Thorn Cactus Tree)**.** *Let* $\widetilde{\mathcal{CT}}(p_1,p_2,p_3,g,w,b)$ *be the set of rooted thorn cactus trees* $\tilde{\tau}$ *consisting of an underlying cactus tree* $\tau \in \mathcal{CT}(p_1,p_2,p_3,g,w,b)$*, and in addition it has:* $n+1-p_1-p_2+g$ *thorns connected to white vertices,* $n-p_2-p_3+w$ *thorns connected to black vertices, and* $n+1-p_1-p_3+b$ thorns *connected to grey vertices. The only restriction for the thorns is:*
(vi) thorns cannot be inside a triangle *of the cactus.*

We are interested in the degree distribution of the white, black, and grey vertices of the cactus trees. Where we define the degree of a vertex v to be:

$$deg(\mathsf{v}) = c + t - \varepsilon + 1, \tag{11}$$

where c is the number of descendants, t is the number of thorns, and ε is 1 if v is the middle vertex of a triangle and is 0 otherwise. *i.e.* it is the standard degree of the vertex, including the thorns but not counting incident non-tree edges, and subtracting one if v is a middle vertex. With this definition of degree, we consider the set $\mathcal{CT}(\lambda,\mu,\nu,g,w,b)$ where $\lambda,\mu,\nu \vdash n$ are the degree distributions of the white, black, and grey vertices respectively. Clearly,

$$\widetilde{\mathcal{CT}}(p_1,p_2,p_3,g,w,b) = \bigcup_{\ell(\lambda)=p_1,\ell(\mu)=p_2,\ell(\nu)=p_3} \widetilde{\mathcal{CT}}(\lambda,\mu,\nu,g,w,b).$$

Example 3. *The thorn cactus tree in Figure 3 belongs to* $\widetilde{\mathcal{CT}}([1^2,2^1,4^1],[1^1,3^1,4^1],[1^1,2^2,3^1],1,1,1)$



Figure 3: Example of a Thorn Cactus Tree

Proposition 2. *The number of thorn cactus trees is:*

$$|\widetilde{\mathcal{CT}}(\lambda,\mu,\nu,g,w,b)| = n\cdot\frac{(\ell(\lambda)-1)!(\ell(\mu)-1)!(\ell(\nu)-1)!}{Aut(\lambda)Aut(\mu)Aut(\nu)}\frac{(g(w-\ell(\nu))+\ell(\mu)\ell(\nu))}{(n-\ell(\lambda)-\ell(\mu)+g+1)}\times$$
$$\times\binom{n-\ell(\lambda)}{w,\ell(\mu)-g-w}\binom{n-\ell(\mu)}{b,\ell(\nu)-w-b}\binom{n-\ell(\nu)}{g,\ell(\lambda)-1-g-b}. \quad (12)$$

The proof of this proposition is postponed to Section 4.

2.3 Reformulation of the main theorem

Let $\mathcal{OP}_r^{(m)}$ be the set of all ordered r-subsets of $[m]$. By definition $|\mathcal{OP}_r^{(m)}| = (m)_r = m(m-1)\cdots(m-r+1)$. We have the following proposition:

Proposition 3.

$$C(\lambda,\mu,\nu) = \sum_{g,w,b\geq 0} |\widetilde{\mathcal{CT}}(\lambda,\mu,\nu,g,w,b)|\ |S_{n+1-\ell(\lambda)-\ell(\nu)+b}|\ |S_{n-\ell(\mu)-\ell(\nu)+w}|\ |\mathcal{OP}^{(n+1-\ell(\lambda)-\ell(\mu)+g)}_{\ell(\nu)-w-b}|. \quad (13)$$

Proof. Note that $C(\lambda,\mu,\nu) = \frac{n!^2}{Aut(\lambda)Aut(\mu)Aut(\nu)}\frac{1}{\binom{n-1}{\ell(\lambda)-1}\binom{n-1}{\ell(\mu)-1}\binom{n-1}{\ell(\nu)-1}}M^{(n-1)}_{\ell(\lambda),\ell(\mu),\ell(\nu)}$. Then, after basic simplifications on the binomial coefficients, a *summand* in the RHS of Equation (13) reduces to:

$$\frac{(n-1)!^2(\ell(\mu)\ell(\nu)+g(w-\ell(\nu)))}{\binom{n-1}{\ell(\lambda)-1}\binom{n-1}{\ell(\mu)-1}\binom{n-1}{\ell(\nu)-1}Aut(\lambda)Aut(\mu)Aut(\nu)}\binom{n}{w,g,b,\ell(\lambda)-1-g-b,\ell(\mu)-g-b,\ell(\nu)-w-b}$$

Then we sum over g, w, and b the terms depending on these parameters. Arranging properly the terms depending on w and b, and simplifying sums on these two parameters thanks to the Vandermonde's convolution formula, we obtain:

$$\sum_{g,w,b} (\ell(\mu)\ell(\nu) + g(w - \ell(\nu))) \binom{n}{w, g, b, \ell(\lambda) - 1 - g - b, \ell(\mu) - g - b, \ell(\nu) - w - b} =$$

$$= n^2 \binom{n-1}{\ell(\nu) - 1} \sum_g \binom{n - \ell(\mu)}{\ell(\lambda) - 1 - g} \binom{n - \ell(\nu)}{g} \binom{n - 1 - g}{n - \ell(\mu)}.$$

Which leads directly to the desired result. □

Combining Proposition 3 and Equation (10), Theorem 1 reduces to:

Theorem 2. *There is an explicit bijection $\Theta^n_{\lambda,\mu,\nu}$ between partitioned 3-cacti in $\mathcal{C}(\lambda, \mu, \nu)$ and tuples $(\tilde{\tau}, \sigma_1, \sigma_2, \chi)$ where $\tilde{\tau} \in \widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$, $\sigma_1 \in S_{n+1-\ell(\lambda)-\ell(\nu)+b}$, $\sigma_2 \in S_{n-\ell(\mu)-\ell(\nu)+w}$, and $\chi \in \mathcal{OP}^{(n+1-\ell(\lambda)-\ell(\mu)+g)}_{\ell(\nu)-w-b}$ for some $g, w, b \geq 0$.*

The next section is devoted to proving this theorem by describing the bijection $\Theta^n_{\lambda,\mu,\nu}$.

3 Description of the bijection

3.1 Additional definitions

Before we get to the description of $\Theta^n_{\lambda,\mu,\nu}$, we need two additional ingredients.

Definition 4 (reverse level traversals (RLT))**.** *For trees $\tilde{\tau} \in \widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$ (or for $\tau \in \mathcal{CT}(\lambda, \mu, \nu, g, w, b)$), we define the white* Reverse Levels Traversal (RLT) *as the traversal going through all the white and black vertices, and all the thorns connected to a white vertex in $\tilde{\tau}$ in the following order:*

(i) the descendants (either black vertices or thorns if any) of the leftmost white vertex of the top white level are traversed from left to right,

(ii) then the leftmost white vertex of the top white level is traversed,

(iii) if i white vertices $(1, \ldots, i)$ (1 being the leftmost) of white level j ($j > 1$) have been traversed and there is a white vertex $i + 1$ at level j on the right of i, the descendants of $i + 1$ are traversed, then vertex $i + 1$ is traversed. Otherwise, the descendants of vertex 1 of white level $j - 1$ are traversed, followed by the white vertex itself.

(iv) The root vertex is the last to be traversed.

The black RLT is defined similarly with respect to black and grey vertices, except that the rightmost black descendant of the white root is the last vertex to be traversed. Similarly, the grey RLT is defined with respect to grey and white vertices, except that the last vertex to be traversed is the rightmost grey descendant of the last black vertex traversed in the black RLT. Figure 3.1 depicts the three RLT for the cactus tree in Example 3.

Definition 5 (Partial permutations)**.** *Given two sets X and Y and a non negative integer m, let $\mathcal{PP}(X, Y, m)$ be the set of bijections from any m-subset of X to any m-subset of Y These bijections are called* partial permutations*. Then $|\mathcal{PP}(X, Y, m)| = \binom{|X|}{m}\binom{|Y|}{m} m!$.*



Figure 4: White, black, and grey RLT on the cactus tree of Example 3

3.2 Bijective mapping Θ for 3-cacti that preserves type

We proceed with the description of Θ:

$$\Theta^n_{\lambda,\mu,\nu} : \mathcal{C}(\lambda,\mu,\nu) \tilde{\rightarrow} \widetilde{\mathcal{CT}}(\lambda,\mu,\nu) \times S_{n+1-\ell(\lambda)-\ell(\nu)+b} \times S_{n-\ell(\mu)-\ell(\nu)+w} \times \mathcal{OP}^{(n+1-\ell(\lambda)-\ell(\mu)+g)}_{\ell(\nu)-w-b}.$$

Within the construction, we use

$$(p_1, p_2, p_3) := (\ell(\lambda), \ell(\mu), \ell(\nu)).$$

3.2.1 The ordered thorn cactus tree $\tilde{\tau}$:

Let $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) \in \mathcal{C}(\lambda,\mu,\nu)$. We construct the ordered tricolored thorn cactus tree $\tilde{\tau} \in \widetilde{\mathcal{CT}}(\lambda,\mu,\nu,g,w,b)$ following the procedure below.

(i) *cactus tree*: The first step is to construct the unlabelled tricolored cactus tree τ and relabelling permutations in the same way as in [17]: Let $\pi_i^{(j)}$ for $i = 1,2,3$ and $j = 1,\ldots,p_i$ be the blocks of the partitions π_i. Also, let $m_2^{'(j)}$ $(1 \le j \le p_2)$ be the maximal element of $\alpha_3^{-1}\alpha_2^{-1}(\pi_2^{(j)})$ and $m_i^{(j)}$ for $i = 1,3$ $(1 \le j \le p_i)$ be the maximal element of the block $\pi_i^{(j)}$. We assign the index p_1 to the block of π_1 containing 1, and suppose that the indexing of the other blocks is arbitrary. We first construct the last passage 3-colored labelled tree T with p_1 white, p_2 black, and p_3 grey vertices satisfying: the root of T is white vertex p_1 and the incidence relations and order of descendants are given in Table 1.

Lemma 1. *The procedure above defines a labelled* 3*-colored tree* T*.*

Proof. Assume that a non root white vertex i_1 is a descendant of grey vertex k, that is in its turn a descendant of black vertex j. Suppose now that j is a descendant of white vertex i_2. Following our construction rules, we have $m_1^{(i_1)} \in \pi_3^{(k)}$, so that $m_1^{(i_1)} \le m_3^{(k)}$. Then $m_3^{(k)} \le m_2^{'(j)}$ as $m_3^{(k)} \in \alpha_3^{-1}(\pi_2^{(j)}) = \alpha_3^{-1} \circ \alpha_2^{-1}(\pi_2^{(j)})$($\pi_2$ is stable by α_2). Finally, as $\alpha_2 \circ \alpha_3(m_2^{'(j)}) \in \pi_1^{(i_2)}$, $\gamma(m_2^{'(j)}) = \alpha_1 \circ \alpha_2 \circ \alpha_3(m_2^{'(j)}) \in \alpha_1(\pi_1^{(i_2)}) = \pi_1^{(i_2)}$ and $\gamma(m_2^{'(j)}) \le m_1^{(i_2)}$. Two cases can occur:

- $m_2^{'(j)} \neq N$ then $\gamma(m_2^{'(j)}) = m_2^{'(j)} + 1 \le m_1^{(i_2)}$. As a direct consequence, $m_1^{(i_1)} < m_1^{(i_2)}$
- $m_2^{'(j)} = N$ and $\gamma(m_2^{'(j)}) = 1 \in \pi_1^{(i_2)}$. As a direct consequence, i_2 is the root vertex.

Incidence relations	Order of descendants
for $1 \le j \le p_1$  if $m'^{(j)}_2 \in \alpha_3^{-1}\alpha_2^{-1}(\pi_1^{(i)})$	 if $\alpha_2\alpha_3(m'^{(j_1)}_2) < \alpha_2\alpha_3(m'^{(j_2)}_2)$
for $1 \le i \le p_3$  if $m_3^{(k)} \in \alpha_3^{-1}(\pi_2^{(j)})$	 if $\alpha_3^{-1}\alpha_2^{-1}\alpha_3(m_3^{(k_1)}) < \alpha_3^{-1}\alpha_2^{-1}\alpha_3(m_3^{(k_2)})$
for $1 \le i \le p_1 - 1$  if $m_1^{(i)} \in \pi_3^{(k)}$	 if $\alpha_3^{-1}(m_1^{(i_1)}) < \alpha_3^{-1}(m_1^{(i_2)})$

Table 1: Incidence relations of tree T.

Rules for adding triangles		
for $1 \le i \le p_1 - 1$, $1 \le j \le p_2$  if $\alpha_2\alpha_3(m'^{(j)}_2) = m_1^{(i)}$	for $1 \le j \le p_2$, $1 \le k \le p_3$  if $\alpha_3^{-1}\alpha_2^{-1}\alpha_3(m_3^{(k)}) = m'^{(j)}_2$ (i.e. $\alpha_3(m_3^{(k)}) = \alpha_2\alpha_3(m'^{(j)}_2)$)	for $1 \le k \le p_3$, $1 \le i \le p_1 - 1$  if $\alpha_3^{-1}(m_1^{(i)}) = m_3^{(k)}$ (i.e. $m_1^{(i)} = \alpha_3(m_3^{(k)})$)

Table 2: Rules for adding non-tree edges to T.

Using a similar argument, one can show that if black vertex j_1 (resp. grey vertex k_1) is on a descending branch of black vertex j_2 (resp. grey vertex k_2), then $m_2^{'(j_1)} < m_2^{'(j_2)}$ (resp. $m_3^{(k_1)} < m_3^{(k_2)}$). Consequently, the white, black, and grey vertices' labels on each branch of the graph are strictly increasing and the last vertex of the branch is always the root vertex (by convention, the block containing 1). ☐

We construct the labelled 3-colored cactus tree Υ from T by adding non-tree edges and forming triangles rooted at the different vertices following the rules in Table 2.

Finally, we remove the labels of Υ to obtain a 3-colored ordered cactus tree τ.

Example 4. *The construction of T, Υ, and τ for the partitioned cactus in Example 2 is depicted in Figure 5.*

(ii) *Relabeling permutations*: These permutations θ_1, θ_2, and θ_3 are defined by considering the *reverse labelled* cactus tree Υ' resulting from the labeling of τ, based on three independent



Figure 5: Tricolored tree and Cactus trees associated to Example 2

reverse-labelling procedure for white, black, and grey vertices. We do a white RLT and label the *white vertices only* (as they are traversed) with the labels $1, 2, \ldots, p_1$. Similarly, we do a black RLT and label the *black vertices only* with labels $1, 2, \ldots, p_2$, and do a grey RLT to label the *grey vertices only* with labels $1, 2, \ldots, p_3$. Next, we relabel the blocks (white, black, and grey) using the new indices from Υ': if a white vertex is labeled i in T and i' in Υ', we set $\pi_1^{i'} = \pi_1^{(i)}$. Black and grey blocks are relabeled in a similar fashion. Let u^i, v^j, w^k be the strings obtained by writing the elements of $\pi_1^i, \pi_2^j, \pi_3^k$ respectively in increasing order. Denote $u = u^1 \cdots u^{p_1}$, $v = v^1 \cdots v^{p_2}$, $w = w^1 \cdots w^{p_3}$ the concatenations of the strings defined above. We define $\theta_1 \in S_n$ by setting u as the first line and $[n]$ as the second line of the two-line representation of this permutation. Similarly, we define the relabeling permutations θ_2 and θ_3 from v and w respectively.

Example 5. *Following up on Example 4, we construct the relabeling permutations θ_1, θ_2, and θ_3. We have:*

$$\theta_1 = \left(\begin{array}{cc|cccc} 4 & 5 & 1 & 2 & 3 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{array}\right), \ \theta_2 = \left(\begin{array}{cccc|cc} 1 & 3 & 4 & 5 & 2 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{array}\right), \ \theta_3 = \left(\begin{array}{c|c|cccc} 5 & 2 & 1 & 3 & 4 & 6 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{array}\right).$$

We define the multisets (allow some elements to be repeated once *i.e.* when there are triangles in Υ'):

$$\begin{aligned} S_1 &= \{\theta_1(m_1^i)\}_{i=1}^{p_1-1} \cup \{\theta_1(\alpha_2\alpha_3(m'^j_2))\}_{j=1}^{p_2} \subset [n], \\ S_2 &= \{\theta_2(m'^j_2)\}_{j=1}^{p_2-1} \cup \{\theta_2(\alpha_3^{-1}\alpha_2^{-1}\alpha_3(m_3^k))\}_{k=1}^{p_3} \subset [n-1], \\ S_3 &= \{\theta_3(m_3^k)\}_{k=1}^{p_3-1} \cup \{\theta_3(\alpha_3^{-1}(m_1^i))\}_{i=1}^{p_1-1} \subset [n-1]. \end{aligned}$$

Note that the sizes of the underlying sets of S_1, S_2, and S_3 are p_1+p_2-1-g, p_2+p_3-1-w, and p_1+p_3-2-b respectively. We use these multisets to double-label the cactus tree τ with three types of labels: *circle* for θ_1, *square* for θ_2, and *rhombus* for θ_3. We assign $(\theta_1(m_1^i), \theta_3(\alpha_3^{-1}(m_1^i)))$ to the white vertices indexed i in Υ', $(\theta_1(\alpha_2\alpha_3(m'^j_2)), \theta_2(m'^j_2))$ to the black vertices indexed j in Υ', and $(\theta_2(\alpha_3^{-1}\alpha_2^{-1}\alpha_3(m_3^k)), \theta_3(m_3^k))$ to the grey vertices indexed k in Υ'. The root vertex is only labeled $n = \theta_1(m_1^{p_1})$.

Let Υ'' be the resulting double-labeled cactus tree. Note that labels (of the same type) of a vertex and its descendant are equal whenever the two vertices belong to a triangle rooted in another vertex.

Let $\overline{S_i}$ (for $i = 1, 2, 3$) be the ordered multiset obtained by arranging the elements of S_i in non-decreasing order (for $i = 1, 2, 3$).

Lemma 2. *Let* $\mathbf{d} = (d_1, d_2, \ldots, d_{p_1+p_2-1})$ *(*$\mathbf{d}' = (d'_1, \ldots, d'_{p_2+p_3})$*, and* $\mathbf{d}'' = (d''_1, \ldots, d''_{p_1+p_3-1})$ *resp.) be the ordered set of labels of the first type (second and third type resp.) obtained by traversing* Υ' *up to, but not including the root vertex according to the white RLT (black and grey RLT resp.) defined in section 3. We have:*

$$\mathbf{d} = \overline{S_1}, \quad \mathbf{d}' = \overline{S_2}, \quad \mathbf{d}'' = \overline{S_3}.$$

Proof. Let $\theta_1(m_1^0) = 0$. By construction, if a black vertex with label $\theta_1(\alpha_2\alpha_3(m'^j_2))$ in Υ'' is a descendant of a white vertex with label $\theta_1(m_1^i)$, $1 \le i \le p_1$, then $\alpha_2\alpha_3(m'^j_2) \in \pi_1^i$ and $\alpha_2\alpha_3(m'^j_2) \le m_1^i$. As θ_1 is increasing among the blocks and within them then:

$$\theta_1(m_1^{i-1}) < \theta_1(\alpha_2\alpha_3(m'^j_2)) \le \theta_1(m_1^i),$$

where the latter is an equality when the white vertex i is the middle vertex of a triangle with black vertex j and rooted at the grey ancestor of vertex i. Now, if black vertices j_1 and j_2 (with labels $\theta_1(\alpha_2\alpha_3(m'^{j_1}_2))$ and $\theta_1(\alpha_2\alpha_3(m'^{j_2}_2))$ in Υ'') are both descendants of the same white vertex, and j_1 is to the left of j_2; then by construction we have $\alpha_2\alpha_3(m'^{j_1}_2) < \alpha_2\alpha_3(m'^{j_2}_2)$. Again, since θ_1 is increasing within blocks of π_1 then $\theta_1(\alpha_2\alpha_3(m'^{j_1}_2)) < \theta_1(\alpha_2\alpha_3(m'^{j_2}_2))$. Finally, the white RLT of the circle labels in Υ'' (up to but not including the root) yields $\overline{S_1}$. Similarly, black and grey RLTs of the square and rhombus labels in Υ'' yield $\overline{S_2}$ and $\overline{S_3}$. ☐

(iii) *thorns*: Recall the definition of $\mathbf{d}, \mathbf{d}'$, and $\mathbf{d}''$ in Lemma 2. We add thorns to the white vertices for each missing element of $[n]$ in $\mathbf{d}$, and add thorns to black and grey vertices for each missing element of $[n-1]$ in $\mathbf{d}'$ and $\mathbf{d}''$, respectively. More specifically, we add $n + 1 - p_1 - p_2 + g$ thorns to the white vertices, $n - p_2 - p_3 + w$ thorns to the black vertices, and $n + 1 - p_1 - p_3 + b$ thorns to the grey vertices of Υ'' in the following fashion: We know that the vectors $\mathbf{d}, \mathbf{d}'$, and $\mathbf{d}''$ are weakly increasing subsequences obtained by the RLTs of Υ''. Below, we describe the rules for adding thorns to the white vertices. By convention, if the vertex is a root of a triangle we do not add thorns *inside* it.

1. If $d_1 > 1$ and the vertex with first label d_1 is white, we connect $d_1 - 1$ thorns to it. If the vertex with first label d_1 is black, we connect $d_1 - 1$ thorns to the ascending white vertex on its left.
2. For $1 < l < p_1 + p_2 - 1$, if $d_l > d_{l-1} + 1$ we follow one of the four following cases:
   (a) d_l and d_{l-1} are both the first label of white vertices in Υ'', white vertex d_l (short for vertex corresponding to d_l) has no black descendant and it is the white vertex following d_{l-1} in the white RLT of Υ''. If so, we connect $d_l - d_{l-1} - 1$ thorns to d_l.
   (b) d_l is the first label of a black vertex and d_{l-1} is the first label of a white one, then black vertex d_l is the leftmost descendant of the white vertex following d_{l-1}. If so, we connect $d_l - d_{l-1} - 1$ thorns to the ascending white vertex of d_l on its left
   (c) d_l is the first label of a white vertex and d_{l-1} is the first label of a black one, then the black vertex d_{l-1} is the rightmost descendant of d_l. If so, we connect $d_l - d_{l-1} - 1$ thorns to d_l on the right of d_{l-1}
   (d) Finally, if d_l and d_{l-1} are both the first label of black vertices, these two vertices have the same white ascending vertex. We connect $d_l - d_{l-1} - 1$ thorns to the ascending white vertex between these two black vertices.

1. if $d_1 > 1$	2. for $1 < l < p_1 + p_2 - 1$, if $d_l > d_{l-1} + 1$			3. if $d_{p_1+p_2-1} < n$
		d_l ○	d_l ●	
	○ d_{l-1}	(a) 	(b) 	
	● d_{l-1}	(c) 	(d) 	

Table 3: Rules for adding thorns to white vertices in Υ''.

3. If $d_{p_1+p_2-1} < n$, we connect $n - d_{p_1+p_2-1} - 1$ thorns to the root vertex on the right of its rightmost black descendant.

We summarize these different cases in Table 3.

Again, we can think of this as adding a thorn to the the white vertices for each element of $[n]$ not included in $\mathbf{d}$.

A similar construction is applied to add thorns to the black and grey vertices following the sequence of integers $\mathbf{d}'$ and $\mathbf{d}''$. Finally we remove all the labels to get the ordered bicolored thorn cactus tree $\tilde{\tau}$

Example 6. *Figure 6 depicts the construction of the thorn cactus tree $\tilde{\tau}$ corresponding to the partitioned cactus in Example 2.*



Figure 6: Construction of the thorn cactus tree associated to Example 2

The next two lemmas show that $\tilde{\tau}$ preserves the type of the partitioned cacti, and that the double-labelled cactus Υ'' can be recovered from $\tilde{\tau}$ via white, black, and grey RLTs.

Lemma 3. *$\tilde{\tau}$ as defined above belongs to $\widetilde{\mathcal{CT}}(\lambda,\mu,\nu,g,w,b)$ where g,w,b are the number of triangles in $\tilde{\tau}$ rooted in grey, white, and black vertices respectively.*

Proof. First we check that the thorns in $\tilde{\tau}$ are not inside the triangles. In the case of the thorns added to the a white vertex i, we only need to consider the instance when i is the middle vertex of a triangle rooted on a grey vertex. This could apply to cases 2. (b) and (c) above. In the former, the thorns cannot be inside a triangle since the non-tree edge would be on the right of the black vertex labeled d_l (recall that in a triangle, the vertices are ordered clockwise as white-black-grey-white). In the latter case, since the label $d_l < d_{l-1}$, then d_l is not the middle vertex of a triangle. A similar conclusion is reached by looking at the rules for adding thorns to black and grey vertices.

Now, we check the vertex degrees of $\tilde{\tau}$. If we take two successive white vertices $i-1$ and i according to white RLT of Υ'' with labels $\theta_1(m_1^{i-1})$ and $\theta_2(m_1^i), (i<p_1)$, a thorn is connected to i for any missing integer of the interval $[\theta_1(m_1^{i-1}), \theta_1(m_1^i)-1]$ in $\mathbf{d}$. This number of missing integers is equal to $\theta_1(m_1^i)-1-\theta_1(m_1^{i-1})-f_i+\varepsilon_i$ where f_i is the number of black descendants of i, and ε_i is 1 if white vertex i is the middle vertex of a triangle and is 0 otherwise. As i is not the root vertex, there is an edge between i and its ancestor so that the resulting degree *deg* for i (as defined in (11)) is:

$$deg(i) = f_i + \big(\theta_1(m_1^i) - 1 - \theta_1(m_1^{i-1}) - f_i + \varepsilon_i\big) - \varepsilon_i + 1 = \theta_1(m_1^i) - \theta_1(m_1^{i-1}),\ \forall i \in [p_1 - 1], \tag{14}$$

Furthermore, $n-\theta_1(m_1^{p_1-1})-f_p$ thorns are connected to the root vertex (since $\varepsilon_{p_1}=0$) so that:

$$deg(p_1) = n - \theta_1(m_1^{p_1-1}) \tag{15}$$

But, according to the construction of θ,

$$\begin{aligned} \theta_1(\pi_1^1) &= [\theta_1(m_1^1)] && (16)\\ \theta_1(\pi_1^i) &= [\theta_1(m_1^i)] \setminus [\theta_1(m_1^{i-1})],\ (2 \le i \le p_1 - 1) && (17)\\ \theta_1(\pi_1^{p_1}) &= [n] \setminus [\theta_1(m_1^{p_1-1})] && (18) \end{aligned}$$

Subsequently:

$$deg(i) = |\pi_1^i|,\ \forall i \in [p_1]. \tag{19}$$

And $\lambda = type(\pi)$ is the white vertex degree distribution of $\tilde{\tau}$. In a similar fashion, μ and ν are the black and grey vertex degree distribution of $\tilde{\tau}$. ☐

Lemma 4. *Assign* circle *labels $1,2,\ldots,n$ to the vertices and the thorns connected to white vertices in $\tilde{\tau}$ in increasing order according to the white RLT, add two other sets of labels $1,2,\ldots,n$ (*square *and* rhombus*) to the vertices and the thorns connected to black and grey vertices in increasing order according to the black and grey RLT. The double-labeling of the vertices is the same as in Υ''.*

Proof. According to the construction of $\tilde{\tau}$, we add thorns to Υ'' when integers are missing in its RLTs so that the thorns would take these missing integers as labels when traversing the thorn cactus tree. As a result, the labels of the vertices in the RLTs of $\tilde{\tau}$ are still $\mathbf{d}, \mathbf{d}'$, and $\mathbf{d}''$ and since they still appear in the same order, we have the desired result. ☐

3.2.2 The ordered set χ and the permutations σ_1 and σ_2

(i) *Ordered set χ:* We define the ordered set $\widetilde{\chi} = \{\theta_1(\alpha_3(m_3^k)) \mid \theta_1(\alpha_3(m_3^k)) \notin S_1\}_{k=1}^{p_3}$. Then, let $\rho : [n] \setminus S_1 \vdash [n - |S_1|]$ be the indexing permutation associating to any integer $i \in [n] \setminus S_1$ its position in $\overline{[n] \setminus S_1}$ where $\overline{[n] \setminus S_1}$ is the ordered (increasing) set of $[n] \setminus S_1$. The ordered set χ is defined as follows:

$$\chi = \rho(\widetilde{\chi}) \tag{20}$$

As $|S_1| = n - (\ell(\lambda) - 1) - \ell(\mu) + g$ and $\left|\{\theta_1(\alpha_3(m_3^k)) | \theta_1(\alpha_3(m_3^k)) \notin S_1\}\right| = \ell(\nu) - w - b$, χ belongs to the set $\mathcal{OP}_{\ell(\nu)-w-b}^{(n+1-\ell(\lambda)-\ell(\mu)+g)}$.

(ii) *Permutations σ_1, σ_2:* Let E and F be the following sets:

$$\begin{aligned} E &= [n] \backslash (\{\theta_1(m_1^i)\}_{i=1}^{p_1-1} \cup \theta_1(\alpha_3(m_3^k))\}_{k=1}^{p_3-1}), \\ F &= [n] \backslash \{\theta_1(\alpha_2\alpha_3(m'^j_2))\}_{j=1}^{p_2} \cup \theta_1(\alpha_3(m_3^k))\}_{k=1}^{p_3-1}). \end{aligned}$$

We define partial permutations $\tilde{\sigma_1}$ and $\tilde{\sigma_2}$ in the following way:

$$\begin{array}{rccc} \tilde{\sigma_1}: & E & \rightarrow & [n-1] \backslash S_3 \\ & u & \mapsto & \theta_3\alpha_3^{-1}\theta_1^{-1}(u) \\ \tilde{\sigma_2}: & F & \rightarrow & [n-1] \backslash S_2 \\ & u & \mapsto & \theta_2\alpha_3^{-1}\alpha_2^{-1}\theta_1^{-1}(u). \end{array}$$

Let $\sigma_1 \in S_{n+1-p_1-p_3+b}$ and $\sigma_2 \in S_{n-p_2-p_3+w}$ be the order isomorphic permutations corresponding to $\tilde{\sigma_1}$ and $\tilde{\sigma_2}$ respectively.

Example 7. *Getting back to Example 2, we have: $\widetilde{\chi} = (4)$ and $\chi = (3)$. Computing the partial permutations leads to:*

$$\tilde{\sigma_1} = \begin{pmatrix} 1 & 3 & 5 \\ 4 & 5 & 3 \end{pmatrix}, \quad \tilde{\sigma_2} = \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix}.$$

Finally:

$$\sigma_1 = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}.$$

3.3 Showing the map Θ is a bijection

To show that $\Theta_{\lambda,\mu,\nu}^n$ is a one-to-one correspondence we take any element $(\tilde{\tau}, \sigma_1, \sigma_2, \chi)$ in

$$(\tilde{\tau}, \sigma_1, \sigma_2, \chi) \in \widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b) \times S_{n+1-\ell(\lambda)-\ell(\nu)+b} \times S_{n-\ell(\mu)-\ell(\nu)+w} \times \mathcal{OP}_{\ell(\nu)-w-b}^{(n+1-\ell(\lambda)-\ell(\mu)+g)}$$

and show that there is a unique element $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2)$ in $\mathcal{C}(\lambda, \mu, \nu)$ such that $\Theta_{\lambda,\mu,\nu}^n(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2) = (\tilde{\tau}, \sigma_1, \sigma_2, \chi)$. Let $p_1 = \ell(\lambda)$, $p_2 = \ell(\mu)$, and $p_3 = \ell(\nu)$. We proceed with a two step proof:

(i) The first step is to notice that $(\tilde{\tau}, \sigma_1, \sigma_2, \chi)$ defines a unique unlabeled cactus tree τ belonging to $\mathcal{CT}(p_1, p_2, p_3, g, w, b)$, a unique multiset $\{S_i\}_{1 \leq i \leq 3}$, as well as a unique ordered set $\widetilde{\chi}$ belonging to $\mathcal{OP}_{p_3-w-b}^{(n+1-p_1-p_2+g)}$. Double labeling each vertex and thorns of τ with $1, 2, \ldots, n$ in increasing order according to the three reverse levels traversals and removing the three sets of thorns (together with their labels) gives a double-labeled cactus tree Υ'' that leads to $\widetilde{\tau}$ according to Θ. This double-labeled cactus tree is the unique one that can lead to $\widetilde{\tau}$ since within Θ, $\widetilde{\tau}$, and Υ'' have the same underlying cactus tree structure τ, and according to Lemma 4, $\widetilde{\tau}$ determines the labels of Υ''.

Then, using Lemma 2, the three series of labels (except the root's) in Υ'' are by construction the three sets $\{S_i\}_{1\leq i\leq 3}$. The knowledge of S_1 and χ uniquely determines $\widetilde{\chi}$. As a result, exactly one 7-tuple $(\tau, S_1, S_2, S_3, \sigma_1, \sigma_2, \widetilde{\chi})$ is associated to $(\tilde{\tau}, \sigma_1, \sigma_2, \chi)$ by the final steps of the mapping Θ.

(ii) The bijection Θ_{n,p_1,p_2,p_3} in [17] is identical to the first steps (up to the construction of $\tau, S_1, S_2, S_3, \sigma_1, \sigma_2$ and $\widetilde{\chi}$) of $\Theta^n_{\lambda,\mu,\nu}$. Therefore there is a unique 5-tuble $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2)$ in $\mathcal{C}(p_1, p_2, p_3, n) = \bigcup_{\ell(\lambda)=p_1,\ell(\mu)=p_2,\ell(\nu)=p_3} \mathcal{C}(\lambda, \mu, \nu)$ mapped to $(\tau, S_1, S_2, S_3, \sigma_1, \sigma_2, \widetilde{\chi})$ by Θ_{n,p_1,p_2,p_3} and equivalently by the first steps of $\Theta^n_{\lambda,\mu,\nu}$.

According to [17], the types of π_1, π_2, and π_3 are directly recovered from $\{S_i\}_{1\leq i\leq 3}$ and τ. Furthermore, using Lemma 3, the vertex degree distribution of $\tilde{\tau}$ is equal to the type of the partitions encoded by the elements in $\{S_i\}_{1\leq i\leq 3}$ corresponding to the relabeling of the maximum elements of the blocks. Finally, as the vertex degree distribution in $\tilde{\tau}$ is (λ, μ, ν), so is the type of (π_1, π_2, π_3). Therefore, $(\pi_1, \pi_2, \pi_3, \alpha_1, \alpha_2)$ belongs to $\mathcal{C}(\lambda, \mu, \nu)$ as desired.

4 Proof of Proposition 2: computation of the number of thorn cactus trees

In this section we prove Proposition 2 where we compute the cardinality of the set $\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$. To do this, we consider its generating function F. The evaluation of F is performed thanks to the multivariate Lagrange inversion theorem (see e.g. [7, 1.2.9]). We propose a recursive decomposition of the desired set of cactus trees sharing similar ideas with [6].

Assume that white, black, and grey vertices are marked respectively by indeterminates x_1, x_2 and x_3. Triangles rooted in a grey, white, and black vertex are marked respectively by x_4, x_5, and x_6. Furthermore, t_i, u_j, and v_k mark respectively white vertices of degree i, black vertices of degree j and grey vertices of degree k. Let $\mathbf{t} = (t_1, t_2, \ldots)$, $\mathbf{u} = (u_1, u_2, \ldots)$, $\mathbf{v} = (v_1, v_2, \ldots)$ and $\mathbf{m}(\epsilon) = (m_1(\epsilon), m_2(\epsilon), \ldots)$ for $\epsilon \vdash n$. As a direct consequence:

$$|\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)| = [x_1^{\ell(\lambda)} x_2^{\ell(\mu)} x_3^{\ell(\nu)} x_4^g x_5^w x_6^b \mathbf{t}^{\mathbf{m}(\lambda)} \mathbf{u}^{\mathbf{m}(\mu)} \mathbf{v}^{\mathbf{m}(\nu)}]\ F \tag{21}$$

In a similar fashion as in [6], we introduce W, B, and G as the generating functions of the sets $\mathcal{W}$, $\mathcal{B}$, and $\mathcal{G}$ of non empty planted thorn cactus trees with respectively white, black, and grey root vertices. Construction rules for these sets of cactus trees are identical to those of $\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$ with the only exception that an additional *planted* edge is connected to the root vertex on the left of the leftmost descendant (vertex or thorn). We take this additional edge into account in the root's degree. Finally, let T_g, T_w, and T_b be respectively the generating functions of triangles rooted in a grey, white, and black vertices. Immediately:

$$T_g = x_4 \tag{22}$$
$$T_w = x_5 \tag{23}$$
$$T_b = x_6 \tag{24}$$

Any cactus tree in $\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$ can be decomposed in a tuple of planted trees in $\mathcal{W}$, $\mathcal{B}$, and $\mathcal{G}$. The rule for the decomposition is based on the nature of the leftmost descendant of the white root in a given cactus tree τ of $\widetilde{\mathcal{CT}}(\lambda, \mu, \nu, g, w, b)$:

(i) If the leftmost descendant is a thorn then τ is equivalent to the cactus tree in $\mathcal{W}$ with the planted edge instead of this leftmost thorn.

(ii) If the leftmost descendant is a black vertex v and v is not the middle vertex of a triangle, then τ is equivalent to the pair (τ_1, τ_2) in $\mathcal{W} \times \mathcal{B}$ where τ_2 is the cactus tree descending from v with v as the root and the edge linking v to the root of τ replaced by the planted edge. τ_1 is the remaining cactus tree descending from the root of τ with the edge linking it to v as the planted edge.

(iii) If the leftmost descendant is a black vertex v_1, and v_1 is the middle vertex of a triangle with grey vertex v_2 then τ is equivalent to the tuple $(\tau_1, \tau_2, \tau_3, t_w)$ in $\mathcal{W} \times \mathcal{B} \times \mathcal{G} \times TT_w$ (TT_w is the singleton set composed of the triangle rooted in a white vertex) where τ_2 and τ_3 are the descending trees from v_1 and v_2 with the edge linking τ's root and v_1 and the edge linking v_1 and v_2 replaced by a planted edge. τ_1 is the remaining descending cactus tree from its root with the leftmost triangle replaced by the planted edge.

One can check easily that the numbers of triangles, white, black, and grey vertices and their degree distribution are stable by the bijective transformation described above. The complicated case above is case (iii) where the edges linking v_1 and v_2, and the edge linking the white root of τ and v_2 are replaced by nothing in (respectively) τ_2 and τ_1. However in the definition 11 of the degree of a vertex in τ, these edges were already not taken into account for the degree of respectively v_1 and the root vertex. As a consequence :

$$F = W + WB + WBGT_w \tag{25}$$

This decomposition is illustrated in Figure 7.



Figure 7: Illustration of the decomposition into planted trees

To determine F, we show that W, B, G, T_g, T_w, and T_b satisfy a system of functional equations. Namely, as shown in Figure 8 any planted cactus tree in $\mathcal{W}$, τ can be decomposed in:

- its white root,
- the cactus trees rooted in a black vertex descending from the root,
- a triple composed of a black rooted cactus tree, a grey rooted cactus tree, a triangle for each triangle descending from the root,
- the positions of the triangles in the descendant list.



Figure 8: Decomposition of a white rooted planted cactus tree

Let i denote the degree of the root vertex (of degree $i+1$), j the number of black descendants not belonging to a triangle and k the number of descending triangles. The vectors $\mathbf{j}$ and $\mathbf{k}$ give the positions of the j black vertices and k triangles within the i descendants. Using the decomposition above, we have:

$$W = x_1 \sum_{i \geq 0} t_{i+1} \sum_{0 \leq j+k \leq i} \sum_{\mathbf{j},\mathbf{k}} B^j (BGT_w)^k \tag{26}$$

Then:

$$\begin{aligned} W &= x_1 \sum_{i \geq 0} t_{i+1} \sum_{0 \leq j+k \leq i} \binom{i}{j, k} B^j (BGT_w)^k \\ W &= x_1 \sum_{i \geq 0} t_{i+1} \left(1 + B + BGT_w\right)^i \\ &= x_1 \Phi_1(W, B, G, T_g, T_w, T_b) \end{aligned} \tag{27}$$

Similarly,

$$\begin{aligned} B &= x_2 \sum_{i \geq 0} u_{i+1} \left(1 + G + GWT_b\right)^i \\ &= x_2 \Phi_2(W, B, G, T_g, T_w, T_b) \end{aligned} \tag{28}$$

$$\begin{aligned} G &= x_3 \sum_{i \geq 0} v_{i+1} \left(1 + W + WBT_g\right)^i \\ &= x_3 \Phi_3(W, B, G, T_g, T_w, T_b) \end{aligned} \tag{29}$$

Finally :

$$(W, B, G, T_g, T_w, T_b) = \mathbf{x} \mathbf{\Phi}(W, B, G, T_g, T_w, T_b) \tag{30}$$

where $\mathbf{x} = (x_1, x_2, x_3, x_4, x_5, x_6)$ and $\mathbf{\Phi} = (\Phi_i)_{1 \leq i \leq 6}$ with $\Phi_i = 1$ for $4 \leq i \leq 6$.

Using the multivariate Lagrange inversion formula for monomials (see [7, 1.2.9]), we find :

$$\begin{aligned} & k_1k_2k_3k_4k_5k_6\,[\mathbf{x}^{\mathbf{k}}]\ W^{r_1}B^{r_2}G^{r_3}T_w^{r_5} = \\ & \qquad \sum_{\{\mu_{ij}\}} \|\ \delta_{ij}k_j - \mu_{ij}\ \| \prod_{1\le i\le 6} [W^{\mu_{i1}}B^{\mu_{i2}}G^{\mu_{i3}}T_g^{\mu_{i4}}T_w^{\mu_{i5}}T_b^{\mu_{i6}}]\Phi_i^{k_i} \end{aligned} \tag{31}$$

where $\mathbf{k} = (k_1, k_2, k_3, k_4, k_5, k_6)$ and the sum is over all 6×6 integer matrices $\{\mu_{ij}\}$ such that:

- $\mu_{11} = \mu_{14} = \mu_{16} = \mu_{22} = \mu_{24} = \mu_{25} = \mu_{33} = \mu_{35} = \mu_{36} = 0$
- $\mu_{ij} = 0$ for $i \ge 4$
- $\mu_{21} + \mu_{31} = k_1 - r_1$
- $\mu_{12} + \mu_{32} = k_2 - r_2$
- $\mu_{13} + \mu_{23} = k_3 - r_3$
- $\mu_{34} = k_4$
- $\mu_{15} = k_5 - r_5$
- $\mu_{26} = k_6$

$$i.e.\ \mu = \begin{bmatrix} 0 & * & * & 0 & * & 0 \\ * & 0 & * & 0 & 0 & 0 \\ * & * & 0 & * & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Looking for zero contribution terms in expression (31), we notice that G and T_w have necessarily the same degree in the formal power series expansion of Φ_1 (see equation (27)). Hence, a non zero contribution of

$$[W^{\mu_{11}}B^{\mu_{12}}G^{\mu_{13}}T_1^{\mu_{14}}T_2^{\mu_{15}}T_3^{\mu_{16}}]\ \Phi_1^{k_1}$$

implies $\mu_{13} = \mu_{15} = k_5 - r_5$. Similar remarks give non zero contributions only for $\mu_{21} = k_6$ and $\mu_{32} = k_4$. As a result, only that one matrix μ yields a non zero contribution.
For this particular μ,

$$\begin{aligned} \frac{1}{k_4k_5k_6}\,\|\ \delta_{ij}k_j - \mu_{ij}\ \| &= r_1\,(k_2k_3 - (k_3 - k_5)k_4) \\ &+ r_2\,(k_6k_3 + (k_6 + r_1 - k_1)(r_3 + k_4 - r_5 - k_3)) \\ &+ r_3\,(k_4k_6 - k_2(k_6 + r_1 - k_1)) \qquad (32) \\ &= \frac{1}{k_4k_5k_6}\Delta(\mathbf{k},\mathbf{r}) \qquad (33) \end{aligned}$$

The next step is to notice that:

$$\begin{aligned} \Phi_1^{k_1} &= \sum_{\mathbf{s}\vdash k_1} \binom{k_1}{\mathbf{s}} \prod_{i\ge 0} \left[t_{i+1}\,(1 + B + BGT_w)^i\right]^{s_i} \\ \Phi_1^{k_1} &= \sum_{\mathbf{s}\vdash k_1} \binom{k_1}{\mathbf{s}} \prod_{i\ge 0} t_{i+1}^{s_i} \sum_{a_1,a_2} \binom{\sum_i is_i}{a_1 - a_2, a_2} B_1^{a}(GT_w)^{a_2} \qquad (34) \end{aligned}$$

As a result, the coefficient in $W^{\mu_{11}}B^{\mu_{12}}G^{\mu_{13}}T_g^{\mu_{14}}T_w^{\mu_{15}}T_b^{\mu_{16}}$ is equal to

$$\sum_{\mathbf{s}\vdash k_1} \binom{k_1}{\mathbf{s}} \prod_{i\ge 0} t_{i+1}^{s_i} \binom{\sum_i is_i}{k_5 - r_5, k_2 - k_4 - k_5 - r_2 + r_5}. \tag{35}$$

Similarly, we have:

$$\begin{aligned} & [W^{\mu_{21}}B^{\mu_{22}}G^{\mu_{23}}T_g^{\mu_{24}}T_w^{\mu_{25}}T_b^{\mu_{26}}]\ \Phi_2^{k_2} = \\ & \qquad \sum_{\mathbf{s}\vdash k_2} \binom{k_2}{\mathbf{s}} \prod_{i\ge 0} u_{i+1}^{s_i} \binom{\sum_i is_i}{k_6, k_3 - k_5 - k_6 - r_3 + r_5} \qquad (36) \\ & \left[W^{\mu_{31}}B^{\mu_{32}}G^{\mu_{33}}T_g^{\mu_{34}}T_w^{\mu_{35}}T_b^{\mu_{36}}\right]\ \Phi_3^{k_3} = \\ & \qquad \sum_{\mathbf{s}\vdash k_2} \binom{k_2}{\mathbf{s}} \prod_{i\ge 0} u_{i+1}^{s_i} \binom{\sum_i is_i}{k_4, k_1 - k_4 - k_6 - r_1} \qquad (37) \end{aligned}$$

Putting everything together gives:

$$
\begin{aligned}
&[x_1^{\ell(\lambda)}x_2^{\ell(\mu)}x_3^{\ell(\nu)}x_4^{g}x_5^{w}x_6^{b}\mathbf{t}^{\mathbf{m}(\lambda)}\mathbf{u}^{\mathbf{m}(\mu)}\mathbf{v}^{\mathbf{m}(\nu)}]\ W^{r_1}B^{r_2}G^{r_3}T_w^{r_5} = \\
&\qquad \frac{\Delta(\ell(\lambda),\ell(\mu),\ell(\nu),g,w,b,\mathbf{r})}{\ell(\lambda)\ell(\mu)\ell(\nu)}\binom{\ell(\lambda)}{\mathbf{m}(\lambda)}\binom{\sum_i i.m_{i+1}(\lambda)}{w-r_5,\ell(\mu)-g-w-r_2+r_5}\times \\
&\qquad\qquad \times\binom{\ell(\mu)}{\mathbf{m}(\mu)}\binom{\sum_i i.m_{i+1}(\mu)}{b,\ell(\nu)-w-b-r_3+r_5} \\
&\qquad\qquad \times\binom{\ell(\nu)}{\mathbf{m}(\nu)}\binom{\sum_i i.m_{i+1}(\nu)}{g,\ell(\lambda)-g-b-r_1}.
\end{aligned}
\tag{38}
$$

Noticing that for $\epsilon \vdash n$

$$
\sum_{i\geq 0} im_{i+1}(\epsilon) = \sum_{i\geq 0}(i+1)m_{i+1}(\epsilon) - \sum_{i\geq 0} m_{i+1}(\epsilon) = n - \ell(\epsilon) \tag{39}
$$

And summing for $\mathbf{r} \in \{(1,0,0,0),(1,1,0,0),(1,1,1,1)\}$ gives the desired result.

5 Proof of Corollary 1 and restriction of bijection Θ when $\nu = [1^n]$

We look more closely at the case when one of the partitions, say ν, is $[1^n]$. We need the following definitions:

Definition 6 (Partitioned bicolored map)**.** *Given partitions $\lambda,\mu \vdash n$, let $\mathcal{C}(\lambda,\mu)$ be the set of triples (π_1,π_2,α) such that $\alpha\in S_n$, π_1,π_2 are set partitions of $[n]$ with type$(\pi_1)=\lambda$ and type$(\pi_2)=\mu$, and each block of π_1 and π_2 is a union of cycles of α and $\beta=\alpha^{-1}\gamma_n$ respectively. The elements of $\mathcal{C}(\lambda,\mu)$ are called* unicellular partitioned bicolored maps *of type λ and μ. Let $C(\lambda,\mu)=|\mathcal{C}(\lambda,\mu)|$.*

Definition 7 (Ordered rooted bicolored thorn trees)**.** *For $\lambda,\mu\vdash n$ such that $\ell(\lambda)+\ell(\mu)\leq n+1$, we define $\widetilde{\mathcal{BT}}(\lambda,\mu)$ as the set of ordered rooted bicolored trees with $\ell(\lambda)$ white vertices, $\ell(\mu)$ black vertices, $n+1-\ell(\lambda)-\ell(\mu)$ thorns connected to the black vertices and $n+1-\ell(\lambda)-\ell(\mu)$ thorns connected to the white vertices. The white (respectively black) vertices' degree distribution (accounting the thorns) is specified by λ (respectively μ). The root is a white vertex.*

Again, adapting the Lagrange inversion developed in [6], we get:

$$
|\widetilde{BT}(\lambda,\mu)| = \frac{n}{Aut(\lambda)Aut(\mu)}\frac{(n-\ell(\lambda))!(n-\ell(\mu))!}{(n+1-\ell(\lambda)-\ell(\mu))!^{\,2}}.
$$

We now prove Corollary 1:

Proof. Since the blocks of π_3 refine the cycles of α_3, if $\nu=[1^n]$ then $\pi_3=\{\{1\},\{2\},\dots,\{n\}\}$ and $\alpha_3=\iota$, the identity permutation. So $\mathcal{C}(\lambda,\mu,[1^n])=\mathcal{C}(\lambda,\mu)$, the number of unicellular partitioned bicolored maps of type λ and μ. Then

$$
\begin{aligned}
\sum_{\lambda,\mu\vdash n} C(\lambda,\mu)m_\lambda(\mathbf{x})m_\mu(\mathbf{y}) &= [m_{1^n}(\mathbf{z})]\sum_{\lambda,\mu,\nu\vdash n} k^n_{\lambda,\mu,\nu}p_\lambda(\mathbf{x})p_\mu(\mathbf{y})p_\nu(\mathbf{z}) \\
&= \sum_{\nu\vdash n,\preceq 1^n\vdash n} R_{\nu,1^n}\sum_{\lambda,\mu\vdash n} k^n_{\lambda,\mu,\nu}p_\lambda(\mathbf{x})p_\mu(\mathbf{y}) \\
&= \sum_{\lambda,\mu\vdash n} k^n_{\lambda,\mu,1^n}p_\lambda(\mathbf{x})p_\mu(\mathbf{y}),
\end{aligned}
$$



Table 4: Local rules for reducing cactus tree $\tilde{\tau}$ when $\nu = [1^n]$.

and $k^n_{\lambda,\mu,1^n} = k^n_{\lambda,\mu}$. By the main theorem, the LHS of Equation (2) is the coefficient of $m_{1^n}(\mathbf{z})$ of the RHS in Equation (1). Since the parameter $g \leq n - \ell(\nu) = 0$, then after some straightforward cancellations we get the desired RHS. ☐

Next, we say what the bijection Θ of Theorem 2 does in this case ($\nu = [1^n]$). This matches the bijection in [14] which in turn matches the bijection in [6] when $\mathsf{g}(\lambda, \mu) = 0$ (there is an analogue of Remark 3 when $\ell(\lambda) + \ell(\mu) = n + 1$).

Corollary 2. *There is a bijection between partitioned bicolored maps* $\mathcal{C}(\lambda, \mu, n)$ *and pairs* $(\tilde{t}, \sigma)$ *where* $\tilde{t} \in \widetilde{BT}(\lambda, \mu)$ *and* $\sigma \in S_{n+1-\ell(\lambda)-\ell(\mu)}$.

Proof. From above we have that $\mathcal{C}(\lambda, \mu, [1^n]) = \mathcal{C}(\lambda, \mu)$. Let $(\tilde{\tau}, \sigma_1, \sigma_2, \chi) := \Theta(\pi_1, \pi_2, [1^n], \alpha_1, \alpha_1^{-1}\gamma)$ for $(\pi_1, \pi_2, \alpha_1) \in \mathcal{C}(\lambda, \mu)$. We know that $\ell(\mu) = n$ forces $\alpha_3 = \iota$, the identity permutation. In this case $m'^{(j)}_2$ is the maximal element of $\alpha_2^{-1}(\pi_2^{(j)})$. But α_2 preserves the blocks of π_2, thus $m'^{(j)}_2$ is just the maximal element of $\pi_2^{(j)}$, call this $m_2^{(j)}$. First, we show that in this case $\tilde{\tau}$ can be reduced to a tree $\tilde{t} \in \widetilde{\mathcal{BT}}(\lambda, \mu)$. Then, we show that σ_1, σ_2 are trivial permutations and that χ can be regarded as a permutation in $S_{n+1-\ell(\lambda)-\ell(\mu)}$.

From the incidence rules in Table 1, we see that each black vertex j has $|\pi_2^j|$ descendants (one for each element of the block). And $\ell(\lambda) - 1$ of the grey vertices have one descendant (one for each $m_1^{(i)}$, $1 \leq i \leq \ell(\lambda) - 1$), the other grey vertices have none. Recall w, b, g count the number of triangles in $\tilde{\tau}$ rooted in white, black, and grey vertices respectively. From the rules in Table (2) for adding triangles rooted at the different vertices, we see that $w = \ell(\mu)$. And if a grey vertex has a white descendant, then these two vertices are part of a triangle rooted in a black vertex, so $b = \ell(\lambda) - 1$. For triangles rooted in grey vertices, if $\alpha_2(m_2^{(j)}) = m_1^{(i)}$ for some i and j ($1 \leq i \leq \ell(\lambda) - 1$ and $1 \leq j \leq \ell(\mu)$), then $m_1^{(i)} \in \pi_2^{(j)}$ and $m_1^{(i)} \leq m_2^{(j)}$ (α_2 preserves blocks of π_2). But $\alpha_1(m_1^{(i)}) \leq m_1^{(i)}$ (α_1 preserves blocks of π_1), so $\gamma(m_2^{(j)}) \leq m_2^{(j)}$. This only happens if $m_2^{(j)} = n$ which means $1 = \gamma(n) \in \pi^{(i)}$ and $i = \ell(\lambda)$, a contradiction. Thus $g = 0$; there are no triangles rooted in a grey vertex.

In terms of the thorns, the cactus $\tilde{\tau}$ has $n + 1 - \ell(\lambda) - \ell(\mu)$ thorns connected to white vertices and since $n - \ell(\mu) - \ell(\nu) + w = 0$ and $n + 1 - \ell(\lambda) - \ell(\nu) + b = 0$, $\tilde{\tau}$ has no thorns connected to black and grey vertices.

From above we see that each grey vertex is either: (i) a middle vertex of a triangle rooted in a black vertex, (ii) a vertex of a triangle rooted in a white vertex, and (iii) a leaf (note that there are $n - (\ell(\lambda) - 1) - \ell(\mu)$ of these). Then depending on the case we do the following reductions: (i) triangle by the non-tree edge, (ii) triangle by the edge between the white root and the black middle vertex, and (iii) leaf by thorn connected to a black vertex. We summarize this reduction graphically in Table 4:

The outcome is an ordered bicolored tree $\tilde{t}$ with $\ell(\lambda)$ white vertices and $\ell(\mu)$ black vertices. This tree $\tilde{t}$ has $n + 1 - \ell(\lambda) - \ell(\mu)$ thorns connected to white vertices and $n + 1 - \ell(\lambda) - \ell(\mu)$ thorns connected to black vertices. Moreover, this reduction $\tilde{\tau} \to \tilde{t}$ is reversible.

In addition, since $\tilde{\tau}$ had no thorns connected to black and grey vertices ($n+1-\ell(\lambda)-\ell(\nu)+b=0$ and $n-\ell(\mu)-\ell(\nu)+w=0$), then σ_1 and σ_2 are trivial permutations. Since $\ell(\nu)-w-b=n+1-\ell(\lambda)-\ell(\mu)=n+1-\ell(\lambda)-\ell(\mu)+g$, then we see that χ is just a permutation σ in $S_{n+1-\ell(\lambda)-\ell(\mu)}$.

In summary, we have a bijection from (λ,μ,n) to the desired pair $(\tilde{t},\sigma)$. ☐

Example 8. *As an example, let* $\alpha_1=(189\,10)(25)(3467)$, $\alpha_2=(15427)(3)(6)(8)(9)(10), \alpha_3=\iota$ ($\alpha_1\alpha_2=\gamma_{10}$)*; and* $\pi_1=\{\{3,4,6,7\},\{1,2,5,8,9,10\}\}$, $\pi_2=\{\{1,2,4,5,7,10\},\{3,9\},\{6,8\}\}, \pi_3=\{\{1\},\{2\},\ldots,\{10\}\}$. *Then* $\Theta(\pi_1,\pi_2,\pi_3,\alpha_1,\alpha_2)=(\tilde{\tau},\emptyset,\emptyset,251364)$ *where* $\tilde{\tau}$ *and its reduction* $\tilde{t}$ *are depicted below:*



Figure 9: Example of restriction of thorn cactus tree to a bicolored thorn trees when $\nu=[1^{10}]$.

References

[1] F. Bédard and A. Goupil The poset of conjugacy classes and decomposition of products in the symmetric group, *Can. Math. Bull*, 35(2):152–160, 1992.

[2] O. Bernardi. An analogue of the Harer-Zagier formula for unicellular maps on general surfaces, *arXiv:1011.2311*. 2010.

[3] P. Biane. Nombre de factorisations d'un grand cycle, *Séminaire Lotharingien de Combinatoire*, 51:4, 2004.

[4] G. Chapuy. *Combinatoire bijective des cartes de genre supérieur*, PhD. thesis, École Polytechnique, June 2009.

[5] G. Chapuy. A new combinatorial identity for unicellular maps, via a direct bijective approach, *DMTCS Proceedings (FPSAC)*, AK:289–300, 2009.

[6] I. P. Goulden and D. M. Jackson. The Combinatorial Relationship Between Trees, Cacti and Certain Connection Coefficients for the Symmetric Group, *European Journal of Combinatorics*, 13:357–365, 1992.

[7] I. P. Goulden and D. M. Jackson. *Combinatorial Enumeration*, Dover, Mineola N.Y. 2004.

[8] I. P. Goulden and A. Nica A direct bijection for the Harer-Zagier formula, *J. Comb. Theory Ser. A*, 111(2):224–238, 2005.

[9] A. Goupil A. and G. Schaeffer. Factoring n-cycles and counting maps of given genus, *European Journal of Combinatorics*, 19:819–834(16), 1998.

[10] J. Irving. On the number of factorizations of a full cycle, *J. Comb. Theory Ser. A*, 113(7):1549–1554, 2006.

[11] D. M. Jackson. Some combinatorial problems associated with products of conjugacy classes of the symmetric group, *J. Comb. Theory Ser. A*, 49(2):363–369, 1988.

[12] S. Lando and A.K. Zvonkin *Graphs on surfaces and their applications,* Springer-Verlag, 2004.

[13] B. Lass Démonstration combinatoire de la formule de Harer-Zagier, *C.R. Acad. Sci. Paris*, 333.Série I:155–160, 2001.

[14] A. H. Morales and E. Vassilieva, Bijective Enumeration of Bicolored Maps of Given Vertex Degree Distribution, *DMTCS Proceedings (FPSAC)*, AK:661–672, 2009.

[15] D. Poulalhon and G. Schaeffer. Factorizations of large cycles in the symmetric group, *Discrete Math.*, 254:433–458, 2000.

[16] G. Schaeffer and E. Vassilieva. A bijective proof of Jackson's formula for the number of factorizations of a cycle, *J. Comb. Theory Ser. A*, 115(6):903–924, 2008.

[17] G. Schaeffer and E. Vassilieva. Partitioned Cacti: a Bijective Approach to the Cycle Factorization Problem, *DMTCS Proceedings (FPSAC)*, AJ:641–652, 2008.

[18] R. P. Stanley. *Enumerative Combinatorics*, volume 2. Cambridge University Press, Cambridge, 1999.

Alejandro H. Morales
Department of Mathematics, Massachusetts Institute of Technology
Cambridge, MA USA 02139
`ahmorales@math.mit.edu`

Ekaterina A. Vassilieva
Laboratoire d'Informatique de l'Ecole Polytechnique
91128 Palaiseau Cedex, France
`katya@lix.polytechnique.fr`